

Evolution of low-mass binaries with black-hole components

L. Yungelson^{a,b,*}, J.-P. Lasota^{b,c}

^a*Institute of Astronomy of the Russian Academy of Sciences, 48 Pyatniskaya Str., 119017 Moscow, Russia*

^b*Institut d'Astrophysique de Paris, UMR 7095 CNRS, Université Pierre et Marie Curie, 98bis Bd Arago, 75014 Paris, France*

^c*Astronomical Observatory, Jagiellonian University, ul. Orla 171, 30-244 Kraków, Poland*

Abstract

We consider evolutionary models for the population of short-period ($P_{\text{orb}} \lesssim 10$ hr) low-mass black-hole binaries (LMBHB) and compare them with observations of soft X-ray transients (SXT). Evolution of LMBHB is determined by nuclear evolution of the donors and/or orbital angular momentum loss due to magnetic braking by the stellar wind of the donors and gravitational wave radiation. We show that the absence of observed stable luminous LMBHB implies that upon RLOF by the low-mass donor angular momentum losses are substantially reduced with respect to the Verbunt & Zwaan “standard” prescription for magnetic braking. Under this assumption masses and effective temperatures (as inferred from their spectra) of the observed donors in LMBHB. Theoretical mass-transfer rates in SXTs are consistent with the observed ones if one assumes that accretion discs in these systems are truncated (“leaky”). We find that the population of short-period SXT is formed mainly by systems which had unevolved or slightly evolved ($X_c \gtrsim 0.35$) donors at the Roche-lobe overflow. Longer period ($P_{\text{orb}} \simeq (0.5 - 1)$ day) SXT might descend from systems with initial donor mass about $1 M_{\odot}$ and $X_c \lesssim 0.35$. It is unnecessary to invoke donors with almost hydrogen-depleted cores to explain the origin of LMBHB. Our models suggest that a very high efficiency of common-envelopes ejection is necessary to form LMBHB, unless currently commonly accepted empirical estimates of mass-loss rates by winds for pre-WR and WR-stars are significantly over-evaluated.

Key words: Binaries: close, stars: evolution, X-rays: binaries

PACS: , 97.10.Cv, 97.10.Gz, 97.80.Jp

1. Introduction

Out of twenty dynamically-confirmed black-hole candidate X-ray binaries ten have K/M spectral type secondaries and orbital periods $\lesssim 1$ day (Remillard and McClintock, 2006). All these systems are transient in X-ray. Their quiescent X-ray luminosity is $\simeq (3 \cdot 10^{30} - 3 \cdot 10^{33}) \text{ erg s}^{-1}$, while in the outburst luminosity may be by a factor $\sim 10^5 - 10^8$ higher. These low-mass black-hole bi-

nary (LMBHB) X-ray systems belong to the class of “Soft X-ray transients” (SXT). Recent reviews of observational data on SXT may be found e.g. in Charles and Coe (2006); Remillard and McClintock (2006). From the evolutionary stand-point SXT are considered as semidetached binaries in which matter is transferred to the compact object via accretion disc. The commonly accepted model for the variability of SXT is based on the thermal-viscous instability of irradiated accretion discs (see Lasota, 2001, and references therein). The estimated number of SXT in the Galaxy ranges from several hundred (Chen et al., 1997) to several thousand (Romani, 1998). Below, we will describe the models

* Corresponding author.

Email addresses: lry@inasan.ru (L. Yungelson), lasota@iap.fr (J.-P. Lasota).




Fig. 1. Initial – final mass relations for massive stars in binaries. The mass of stars at terminal main-sequence (TAMS), the initial mass of helium core $M_{\text{core},i}$, the pre-supernova mass $M_{\text{pre-SN}}$, and the black-hole mass M_{BH} are shown.

of a population of short-period LMBHB, discuss their observational properties, and consider possible existence of a population of faint LMBHB with quasi-stable, cold accretion discs (Menou et al., 1999). This review is based mainly on the studies by Yungelson et al. (2006) (henceforth, Paper I) and Yungelson & Lasota (in prep., Paper II).

2. The model

2.1. Formation of black holes with low-mass companions

Modeling of the Galactic population of LMBHB includes two major steps: (a) modeling of the population of zero-age binaries containing a black hole accompanied by a low-mass companion and (b) following the evolution of the system to Hubble time.

The population of zero-age LMBHB was computed (Paper I) using population synthesis code SEBA

(Portegies Zwart and Verbunt, 1996; Portegies Zwart and Yungelson,

1998; Nelemans et al., 2001; Nelemans et al., 2004) with 250000 initial binaries with $M_{10} \geq 25 M_{\odot}$. Assumed time- and position-dependent Galactic star formation history is based on the model of Boissier and Prantzos (1999), but for the inner 3 kpc of the Galaxy we doubled the star formation rate given in the latter study to mimic Galactic bulge. We assumed a 50% binarity rate (2/3 of stars in binaries), an IMF after Kroupa et al. (1993), an initial distribution of separations in binaries (a) flat in $\log a$ between contact and $10^6 R_{\odot}$, a flat mass ratio distribution, and initial distribution of

eccentricities of orbits $\Xi(e) = 2e$.

A summary of the assumptions on transformations of stellar masses in the course of the evolution is shown in Fig. 1. Black hole progenitors have $M_0 = 25 - 100 M_{\odot}$. The algorithm for formation of black holes in the code follows the fall-back scenario (Fryer and Kalogera, 2001) under assumption that explosion energy is fixed at 10^{50} ergs, which is within the expected range but favours formation of rather massive black holes (up to $\sim 15 M_{\odot}$). Nascent black holes receive an asymmetric kick at formation following Paczyński (1990) velocity distribution with $\sigma_v = 300 \text{ km s}^{-1}$, but scaled down with the ratio of the black hole mass to a neutron star mass. The population of LMBHB is not sensitive to the assumed kick distribution, since scaled down with $M_{\text{bh}}/M_{\text{ns}}$ -ratio kick velocities are too small to disrupt close binaries in SN explosions. Test run for Maxwellian kick distribution for pulsars with $\sigma = 265 \text{ km/s}$ (Hobbs et al., 2005) confirmed this assumption. For the common-envelope phase we used the standard prescription (Webbink, 1984; de Kool et al., 1987), with the efficiency and structure parameters α_{CE} and λ product $\alpha_{\text{CE}} \times \lambda = 2$ (see discussion of the issue of $\alpha_{\text{CE}} \times \lambda$ in §5). For more detailed description of input parameters we refer the reader to Paper I.

2.2. Angular momentum loss

Since low-mass components in LMBHB have KV/MV spectral types, it is usually assumed that evolution of LMBHB is governed by angular momentum losses (AML), like for cataclysmic variables. In our computations we assumed that AML via magnetic stellar wind (MSW) follows the model suggested by Verbunt and Zwaan (1981):

$$\frac{d \ln J}{dt} = -\frac{1}{2} \times 10^{-28} k^2 \left(\frac{2\pi}{P} \right)^{10/3} \frac{1}{G^{2/3}} \frac{M_t^{1/3} R_2^4}{M_1 f^2}, \quad (1)$$

where M_1 is mass of the primary, M_t – total mass of the system, P – orbital period, R – radius of the secondary, $k^2 \sim 0.1$ – its gyration radius, $f \sim 1$ – a parameter derived from observations. Note, that this model extrapolates via almost two orders in magnitude stellar rotation braking law derived by Skumanich (1972) for *single* field stars to rapidly rotating components of close binaries and assumes efficient spin-orbital coupling.

Angular momentum loss via gravitational waves radiation (GWR) is described by the standard Landau and Lifshitz (1971) formula




Fig. 2. Fractions of low-mass bh+ms binaries formed at different epochs in the lifetime of the Galaxy (solid line) and the shape of star formation rate (SFR) adopted in present study (dashed line). At maximum, the SFR is $\simeq 15.6 \text{ M}_{\odot} \text{ yr}^{-1}$. Current SFR is $\simeq 1.9 \text{ M}_{\odot} \text{ yr}^{-1}$.

$$\frac{d \ln J}{dt} = -\frac{32}{5} \frac{G^{5/3}}{c^5} \left(\frac{2\pi}{P} \right)^{8/3} \frac{M_1 M_2}{(M_1 + M_2)^{1/3}}. \quad (2)$$

Below, we will refer to the model in which semidetached systems obey Eqs. (1) and (2) as to a “standard” model.

2.3. Population of unevolved LMBHB

With our assumptions we find that, within a Hubble time (13.5 Gyr), $\sim 49\,000$ binaries that have orbital periods below 2.0 day and contain black holes accompanied by main-sequence stars with $M_{20} < 1.6 \text{ M}_{\odot}$ (for which magnetic braking is supposed to operate) were formed in the Galaxy.¹ Out of them, $\sim 17\,000$ were brought into contact by AML via MSW and/or GWR and $\sim 14\,000$ evolved to shorter periods under “standard” assumptions on MSW (see § 2.2).

Because black holes form in the first ~ 3 Myr after their progenitor formation, formation history of bh+ms binaries strictly follows the star formation history (Fig. 2). Figure 3 shows relations between initial masses of components in model systems and their post-circularization orbital period². Solid lines in the upper panel of the Figure outline the

¹ This number is $\sim 0.01\%$ of all binaries with black-hole components formed in Hubble time.

² Every dot in the plot actually represents four or five systems with similar combination of M_{10} , M_{20} , P_0 , but born at different time. Since we present one random realization of the model, all numbers given are subject to Poisson noise.




Fig. 3. Upper panel: relation between orbital periods and masses of main-sequence companions of black holes after circularization of the orbits. Systems with main-sequence components that may fill their Roche lobes in their main-sequence lifetime or in the Galactic disc lifetime (13.5 Gyr) are located to the left of the solid curves that correspond to $M_{\text{BH}} = 4 \text{ M}_{\odot}$ (left one) and $12 \text{ M}_{\odot}$ (right one), respectively. Lower panel: relation between initial masses of components in bh+ms systems.

mass and period range from which systems may be brought in the contact via action of AML, however, in “standard” model only systems that overflow Roche lobes at periods less than $\simeq 15$ hr evolve to shorter periods (the rest have small helium cores and evolve to longer periods, forming at the end helium white dwarfs, if time span between contact and Hubble time is large enough; otherwise, at present their donors are subgiants).

3. Evolution of LMBHB

For simulation of the current Galactic population of semidetached LMBHB we convolved the above-described “underlying” population of binaries born at different epochs in the history of the Galaxy with the grids of evolutionary tracks for low-mass components in the binaries with different combinations of masses of components and post-circularization (initial) orbital periods. For

evolutionary computations, appropriately modified TWIN version (September 2003) of Eggleton (1971) evolutionary code was used. Every “initial” system was “evolved” through $13.5 \text{ Gyr} - T_0$, where T_0 is the moment of formation. In the Figures presented below, we show only systems that have at present $q = M_2/M_1 \geq 0.02$. At $q \lesssim 0.02$ the circularization radius of accretion stream becomes greater than the outer radius of the accretion disc, resonance phenomena in the disc become important and, though computations may be continued in the same fashion as before, until models cease to converge due to very low mass, actually it is unclear how mass transfer occurs (see Paper I for more details). The systems with $q < 0.02$ have $P_{\text{orb}} \lesssim 2 \text{ hr}$ and mass-transfer rates ($< 10^{-10} M_{\odot} \text{ yr}^{-1}$). In the “standard” model currently more than 75% of all low-mass ms+bh systems have $q < 0.02$ and have yet to be observed if mass-transfer occurs in them.

3.1. Luminous persistent LMBHB?

Figure 4 shows the distribution of systems with $q > 0.02$ in the “standard” model over mass-transfer rates. The upper two panels show the break-down of the binaries that have stable or unstable accretion discs (370 and 2900 objects, respectively). Irradiated discs are hot and stable if accretion rate (in $M_{\odot} \text{ yr}^{-1}$) exceeds (Dubus et al., 1999)

$$\dot{M}_{\text{crit}}^+ \approx 2.4 \times 10^{-11} M_{\text{BH}}^{-0.4} R_{d,10}^{2.1} \left(\frac{\mathcal{C}}{5 \times 10^{-4}} \right)^{-0.5}, \quad (3)$$

where M_{BH} is the mass of the accretor in $M_{\odot}$, $R_{d,10}$ is the disc outer radius in 10^{10} cm , and $\mathcal{C}$ is a measure of heating of the disc by X-rays (Shakura and Sunyaev, 1973); for a given disc radius R the irradiation temperature T_{irr} is

$$T_{\text{irr}}^4 = \mathcal{C} \frac{\dot{M} c^2}{4\pi\sigma R^2}, \quad (4)$$

where σ is the Stefan-Boltzmann constant. We use $\mathcal{C} = 5 \times 10^{-4}$, following Dubus et al. (1999) who have found that this value is consistent with properties of persistent low-mass X-ray sources, but actually $\mathcal{C}$ might e.g. vary in time (Esin et al., 2000).

However, no persistent LMBHB with $P_{\text{orb}} \lesssim 4 \text{ hr}$ are observed³. Stable model systems have an average luminosity $L = 0.1 \dot{M} c^2 \approx 5 \cdot 10^{36} \text{ erg s}^{-1}$. The accretion luminosity may be assumed to be emitted

³ In fact no LMBHB at all are observed below $P_{\text{orb}} \sim 4 \text{ hr}$.




Fig. 4. Mass transfer rates vs. orbital period for the model population of semidetached low-mass binaries with assumed black hole accretors. Upper panel – stable systems in the model for a “standard” Verbunt-Zwaan assumption on AML via MSW. Middle panel – population of unstable systems in the same model. Lower panel – population for the model in which MSW does not operate in semidetached systems. In this model all binaries have unstable discs. The arrows mark approximate upper limits to the average mass-transfer rates for SXT under assumption of “leaky” discs, while plusses show estimates of rates based on assumed recurrence period of 30 years, except for V616 Mon and IL Lup whose recurrence times are known to be ~ 50 and 10 years, respectively (see § 3.3.2 for details).

as that of a 1 keV blackbody, corresponding to the “high/soft” X-ray spectral state of LMBHB. Given their location in the Galaxy provided by the population synthesis code, we may roughly estimate intervening X-ray absorption. We found that the number of persistent systems expected above the completeness limit for the X-ray sources seen by the *RXTE All Sky Monitor* (about 3 mCrab or 0.2 counts/s) in the $2\text{--}10 \text{ keV}$ band is 85. Setting the threshold ten times higher still yields 13 bright, persistent LMBHB with *ASM* fluxes comparable to e.g. SMC X-1. Noticeable number of stable “observed” persistent model sources remain in the model also under another assumptions on the spectrum of their emission (see for details Paper I). If the discs were not irradiated, no systems would be stable. However, both persistent neutron-star low-mass X-ray binaries and outbursting black hole transients show clear signatures of irradiation. Note that with parameter $\mathcal{C} =$

5×10^{-4} , only a 0.5% of the accretion luminosity (efficiency of 10%) is reprocessed in the disc.

Reduction of $d \ln J / dt$ by factor 2 compared to Eq. (1) does not change the results significantly (see Paper I) and points to the necessity of stronger reduction of AML for semidetached systems.

3.2. Reduced angular momentum loss?

The problem of production of unobserved persistent low-mass black-hole X-ray binaries under “standard” assumptions was noted before e.g. by King et al. (1996); Ergma and Fedorova (1998); Menou et al. (1999); Ivanova and Kalogera (2006). Skumanich (1972) “law” on which Verbunt and Zwaan (1981) model is based, is apparently in conflict with observational data on rotation velocities in young open clusters (Collier Cameron, 2002; Andronov et al., 2003). According to the latter study, the time-scale of rotation braking is two orders of magnitude longer than the one based on Skumanich law. Necessity for “weaker” than Verbunt & Zwaan AML by MSW was suggested e.g. by Hameury et al. (1988); Ivanova and Taam (2003) for cataclysmic variables and by van Paradijs (1996) for SXT.

In Paper I we suggested, in line with observational evidence mentioned above, that magnetic braking operates on a much reduced scale (as compared with Verbunt and Zwaan (1981) prescription) or that it does not operate at all in the semidetached systems with black-hole accretors. As a test of this hypothesis, we computed a population of LMBHB under assumption that MSW is not operating once the RLOF occurs.

Since the mechanism that brings the systems to the RLOF is the same both in “standard” and in “no-MSW” cases, the number of semidetached systems is the same in both models ($\simeq 14000$), but in the latter case, because of a weaker AML, there are currently in the Galaxy about 5000 LMBHB that evolve to shorter periods and have $q \geq 0.02$ (compared to $\simeq 3200$ in the former case). All these systems are transient according to the criterion (3). This population is plotted in the lower panel of Fig. 4.

Our estimate of the number of SXT apparently exceeds the estimates based on observations. Assuming that their outbursts peak on average at $0.1 L_{\text{edd}}$ in X-rays, all of these transients should be seen in outburst by the *RXTE ASM*, regardless of their lo-

Table 1
Observed short-period low-mass binaries.

Object	P_{orb} , hr	Sp	q	M_{bh} , $M_{\odot}$
1 XTE J1118+480	4.10	K5V-M1V	0.044 - 0.035	6.5 - 7.2
2 GRO J0422+32	5.09	M2 $\pm$ 2V	0.313 - 0.008	3.7 - 5.0
3 GRS 1009-45	6.84	G5V-M0V	0.159 - 0.125	3.6 - 4.7
4 XTE J1650-500	7.68	K4 V	0.1	3.4 - 7.9
5 A0620-00	7.75	K3V-K7V	0.075 - 0.055	8.7 - 12.9
6 GS 2000+25	8.28	K3V-K6V	0.053 - 0.035	7.2 - 7.8
7 XTE J1859+226	9.12	G5V-K0V		
8 GRS 1124-68	10.39	K3V-K7V	0.208 - 0.114	6.5 - 8.2
9 H 1705-25	12.50	K3V-K7V	< 0.053	5.6 - 8.3

cation in the Galaxy. Matching the observed discovery rate of SXT would require the outbursts to be sub-Eddington at maximum and/or recurrence times to be $\gtrsim 100$ years for most sources.

In the “no-MSW” case, donor-stars in the LMBHB have $X_c \gtrsim 0.35$ at RLOF, while in the “standard” case all systems which did not exhaust hydrogen in the core to the instant of the RLOF evolve to the shorter P_{orb} . This also means that the range of initial periods of systems in “no MSW” case is smaller: it extends to ≈ 35 hr.

3.3. Observational parameters of short-period LMBHB

In Table 1 we summarized the main parameters of short orbital period SXT that are important from evolutionary point of view. The range of the spectral types assigned to secondaries is based on published data, see for references Paper II; the values of mass ratios of components $q = M_2/M_{\text{bh}}$ and the estimates of black-hole masses are taken from Orosz (2003); Orosz et al. (2004).

3.3.1. Effective temperatures and masses of donors

The essential information available on the companion-stars in SXT is their spectral types. Determination of the latter is a challenging task since the emission of the cool star is contaminated by the radiation from the accretion disc and the hot spot where the accretion stream hits the disc’s edge (Charles and Coe, 2006). Different methods of Sp -determination result in a scatter of assigned spectra that may amount to several subtypes (Table 1). In the absence of direct determinations of such pa-




Fig. 5. Effective temperatures of donors in model population for “no-MSW” case (dots). Vertical lines mark the ranges of T_{eff} of donor-stars in observed SXT corresponding to the ranges of the estimates of their spectral types (Table 1). Systems are annotated according to their numbers in Table 1. Large filled circles give T_{eff} of donors derived from the fits to the synthetic spectra. Dashed lines are evolutionary tracks for $M_0=1.0$ and $1.1 M_\odot$ stars with $4 M_\odot$ companions. Heavy solid lines to the left and right, respectively, show “limiting” tracks for $(1+12) M_\odot$, $P_0 = 9.6$ hr system in which MSW does not operate after RLOF and $(1+4) M_\odot$, $P_0 = 45.6$ hr system with MSW operating after RLOF (see § 3.3.1 for data on tracks and discussion). $Sp - T_{\text{eff}}$ relation used in the paper is shown at the right border of the coordinate box.

rameters of the stars as their effective temperatures and masses (for a few exceptions see below), one is forced to use $Sp - T_{\text{eff}}$ and $Sp - M$ relations. We adopted for the present study $Sp - T_{\text{eff}}$ relation by Tokunaga and $Sp - M$ relation by Schmidt-Kaler given in Cox (2000). In Figs. 5 and 6 we plot, respectively, effective temperatures of the donors and their masses in model population vs. orbital periods of the systems. In the absence of objective criteria for discrimination of reliable vs. non-reliable Sp -determinations, we compare model data with the ranges of T_{eff} and M_2 inferred from the spectra.

Within uncertainties of the Sp -determinations and conversion $Sp - T_{\text{eff}}$ the model satisfactorily reproduces T_{eff} and M_2 of the donors in the LMBHB with $P_{\text{orb}} \lesssim 9$ hr. Several systems – XTE 1650-500, A0620-00, and GS 2000+25 – seem to be located below the “populated” areas of the plots. However, we should note the following. In order to avoid interpolation between pre-computed tracks that evolve upon RLOF continuously to shorter periods and tracks that turn to longer periods immediately or change the direction of evolution in P_{orb} , we restricted ourselves by binaries that evolve to short P_{orb} . For a system with given M_{10} and M_{20} the direction of evolution changes quite abruptly,




Fig. 6. Masses of donor-stars in the modeled population for “no-MSW” case. Vertical lines show the ranges of M_2 corresponding to the range of spectral types of donors in observed SXT (Table 1). Dashed and continuous lines are the same evolutionary tracks as in Fig. 5. For XTE J1650-500 (at $P_{\text{orb}}=7.88$ hr) there is only one determination of the spectrum – K4V, but the same authors mention that the next best fits are G5V and K2III; for this reason we show by solid circle the lower limit for M_2 in this system.

over a narrow range of initial $\Delta P_{\text{orb}} \lesssim 0.1$ day. For a given combination of M_{10} and M_{20} a “gap” between tracks evolving in different directions forms. This is seen from Fig. 5 where we plot evolutionary tracks for $(1+4) M_\odot$, $P_0=1.4$ and 1.5 day and for $(1.1+4) M_\odot$, $P_0=1.3$, 1.4 , and 1.45 day systems. But since there is a continuity in the initial parameters of the systems, the “gap” has to be filled by the systems that start RLOF in the well populated area in the $P_0 - M_{20}$ diagram (Fig. 3). Thus it is at least qualitatively clear that the origin of short-period LMBHB may be explained within paradigm of strong reduction of magnetic braking in systems with donors overflowing Roche lobes.

In our model we reduced AML by MSW to zero. In reality some AML by MSW can be still operating. We plot in Figs. 5 and 6 two “limiting” tracks: for $(1+12) M_\odot$, $P_{\text{orb},0} = 9.6$ hr in which the donor overflows Roche lobe almost unevolved and MSW is absent after RLOF and for $(1+4) M_\odot$, $P_{\text{orb},0} = 45.6$ hr in which donor has $X_c \simeq 10^{-4}$ at RLOF and MSW continues to operate. Crudely, model populations with MSW and without MSW that evolve to short P_{orb} have to be located between these two limiting curves. There is a contribution of lower and higher mass systems of course as we plotted tracks

for $1M_{\odot}$ donors for simplicity. Adding some AML to our model will shift the population to the right, giving a better agreement with observations while still not producing stable luminous X-ray sources. From Fig. 6 in Paper I it is seen that such an adding of AML will influence mainly the long-period systems.

Our computations for the “no-MSW” model show that LMBHB evolve to longer periods if $X_c \lesssim 0.35$ at RLOF. For instance a $1.1M_{\odot}$ donor with $P_{\text{orb}0} = 1.4$ (middle curve) spends in the RLOF-state almost 10 Gyr, out of which about 5 Gyr it evolves to longer periods. This provides a possibility of explaining SXTs with periods > 9 -10 hr.

3.3.2. Mass-transfer rates

Our model predicts the distribution of SXT over mass-transfer rates $\dot{M}$ that can also be compared to $\dot{M}$ deduced from observations of SXT sources.

Usually, $\dot{M}$ for SXT are evaluated by dividing the mass accreted during outburst by the recurrence time (see e.g. White and van Paradijs, 1996). Among short-period SXT the recurrence time is known only for A0620-00 (about 60 years) and 4U 1543-47 (about 10 years). For the other systems one can only obtain upper limits of $\dot{M}$ since only one outburst has been seen in the X-ray observations epoch (30 years are usually assumed). These estimates of $\dot{M}$, derived from the data on outburst parameters from Chen et al. (1997) and for KV UMa and V406 Vul based of observations presented in Chaty et al. (2003) and Hynes et al. (2002) respectively, are shown as plusses in Figs. 4 and 7.

For the “standard” model all above-described $\dot{M}$ -estimates are inconsistent with the model (two upper panels of Fig. 4). Even if only GWR acts, $\dot{M} \sim 10^{-12} M_{\odot} \text{ yr}^{-1}$ cannot be explained if one assumes that they represent *secular* values (lower panel of Fig. 4). It would take post-minimum period systems several Hubble times to reach $P_{\text{orb}} \gtrsim 4$ hr and have a secular $\dot{M} \sim 10^{-12} M_{\odot} \text{ yr}^{-1}$. If the extremely low $\dot{M}$ deduced for short period SXTs were due to downward fluctuations from an intrinsically high secular $\dot{M}$, one would still have to explain why the bright counterparts are not observed (§ 3.1). Furthermore, some systems would then necessarily have higher rates than their secular value, hence there should be even more persistent systems.

In Paper I we have shown that in the “standard” model systems with $P_{\text{orb}} \approx 8 - 10$ hr may be consistent with systems where donors overflow Roche lobes extremely close to TAMS, but then one needs

to assume some very special initial distribution of binaries over orbital separations which will after several evolutionary stages with mass transfer and mass and momentum loss from the system and supernova explosion, result in concentration of zero-age black-hole and main-sequence-star systems just at the desired very narrow range of separations. On the other hand, if a small amount of AML is added to the alternative “no-MSW” model, the latter will become consistent with recurrence time based estimates of $\dot{M}$ for observed $P_{\text{orb}} \approx 8 - 10$ hr systems, as shows comparison of two lower panels of Fig. 4.

The longest period SXT of the “short-period” group, 4U1543-47 ($P_{\text{orb}} = 27$ hr; it is not shown in the Figures) is, in principle, consistent with the evolution of “standard” model systems with donors overflowing Roche-lobes after formation of He-cores. But again, these binaries evolve through the Hertzsprung gap extremely fast, and the probability of observing such systems is very low (see Kolb, 1998). In the “no-MSW” model, formation of the systems similar to 4U1543-47 is much easier explained, since to the range of periods about 13 hr slowly enough evolve binaries with $M_{20} \simeq 1 M_{\odot}$ donors which overfilled Roche lobes having $X_c \lesssim 0.35$.

3.3.3. Truncated discs

It is, however, plausible that the $\dot{M}$ in short-period SXTs are much higher than $\dot{M}$ derived above. These $\dot{M}$ were obtained treating accretion disc as a reservoir which during quiescence is filled up to a critical value at which the outburst is triggered and the disc emptied. The implicit assumption is that the reservoir is not leaky, i.e. that the accretion rate at the disc’s inner edge is much smaller than $\dot{M}$. However, the inner truncation of the discs is required to explain observed quiescent luminosities (e.g. Lasota, 1996; Lasota et al., 1996). The same may be true for non-quiescent discs too (see e.g. Done and Gierliński, 2006). Dubus et al. (2001) showed that the disc instability model can reproduce the observed X-ray light-curves only if discs in SXTs are truncated and irradiated. The truncated (or “leaky”) disc paradigm is now commonly used in describing the observed timing and spectral properties of X-ray LMBHB (McClintock and Remillard, 2006, and references therein).

For systems with *non-stationary* quiescent accretion discs $\dot{M}$ can be estimated as (Lasota, 2001)

$$\dot{M} \approx \frac{\epsilon \dot{M}_{D,\max}}{t_{\text{recc}}} + \dot{M}_{\text{in}} \quad (5)$$

where

$$\dot{M}_{D,\max} = 2.7 \times 10^{21} \alpha^{-0.83} \left(\frac{M_1}{M_\odot} \right)^{-0.38} R_{d,10}^{3.14}, \quad (6)$$

is the maximum quiescent-disc mass (in g) and $\epsilon = \Delta M_D / M_{D,\max}$ – the fraction of the disc’s mass lost during outburst; α is the kinematic viscosity parameter, $R_{d,10}$ is the disc’s outer radius in 10^{10} cm and $\dot{M}_{\text{in}}$ is the accretion rate at the disc’s inner edge. The usual, non-leaky disc estimates, neglect $\dot{M}_{\text{in}}$. However, in SXTs $\dot{M}_{\text{in}}$ is the dominant term as argued by Menou et al. (1999).

According to the DIM, the disc is in cold thermal equilibrium if accretion rate at all annuli satisfies the inequality (Lasota et al., 2008)

$$\begin{aligned} \dot{M}(r) &< \dot{M}_{\text{crit}}^- \\ &\approx 2.64 \times 10^{15} \alpha_{0.1}^{0.01} R_{10}^{2.58} \left(\frac{M_1}{M_\odot} \right)^{-0.85} \text{ g s}^{-1}, \end{aligned} \quad (7)$$

where $\alpha_{0.1}$ – the viscosity parameter in 0.1. Therefore, the mass accretion rate at the truncation radius ($\dot{M}_{\text{in}}$) must be smaller than $\dot{M}_{\text{crit}}^-(r_{\text{in}})$. Parameterizing the truncation radius as a fraction of the circularization radius $r_{\text{in}} = f_t r_{\text{circ}}$ (where $f_t < 0.48$, see Menou et al., 1999) one gets (in $M_\odot \text{ yr}^{-1}$)

$$\begin{aligned} \dot{M}_{\text{max}} &\lesssim 2.5 \times 10^{-7} \\ &\left[(1+q)^{1/3} (0.5 - 0.227 \log q) \right]^{10.32} P_d^{1.72} f_t^{2.58}, \end{aligned} \quad (8)$$

where period is in days. Inserting in Eq. (8) the lower limits on q from Table 1 and assuming $f_t = 0.48$, one obtains the upper limits to the mass-transfer rates shown by arrows in Figs. 4 and 7. One can therefore conclude that a substantially reduced MSW AML would be consistent with both the stability properties and $\dot{M}$ of SXTs.

4. Cold, stable systems

Possible problem of excess of SXT in our model mentioned in § 3.1 may be resolved in the model suggested by Menou et al. (1999) (henceforth, MNL). MNL pointed out that if the truncation radii were slightly larger than estimated to fit observations of quiescent SXTs the discs would be globally stable, since their accretion rates are lower than the critical value given by Eq. (7) corresponding to cold stable equilibria.




Fig. 7. Mass transfer rates vs. orbital period for the model population of semidetached low-mass binaries with assumed black hole accretors. Upper panel – model for Verbunt-Zwaan assumptions on AML via MSW. Crosses mark systems that are stable according to Menou et al. criterion for instability of cold discs, while dots mark unstable systems. Middle panel shows position of systems with discs stable according to Menou et al. criterion in the model in which MSW does not operate in semidetached systems. Lower panel – systems with unstable discs in model without MSW. Arrows and plusses have the same meaning as in Fig. 4.

Figure 7 shows the distribution of stars in two population models with respect to stability criterion suggested by MNL under assumption the truncation occurs at $r_{\text{in}} = 0.48 r_{\text{circ}}$. In the “standard” case with MSW AML all ($\simeq 2900$), but very few ($\simeq 60$), systems are unstable (Fig. 7 upper panel) vs. MNL-criterion. On the other hand, in the model without MSW after RLOF there are almost equal numbers ($\simeq 2500$) of stable and unstable vs. MNL-criterion systems (Fig. 7, two lower panels). This is because $\dot{M}$ in the GWR-only scenario are lower than in the MSW+GWR scenario, thereby enabling the criterion for a cold disc to be met more easily. Truncating the disc at R_{circ} would make all transients “MNL-stable” for both models. Therefore, it is possible that most or even all LMBHB are secularly cold/stable systems, erupting from time to time due to randomly acting factors like variability of truncation radii or mass transfer rate. The predicted population size of these systems is too small to contribute significantly to the large number of faint ($L_X \sim 10^{31} \text{ erg s}^{-1}$) hard X-ray sources seen

in the deep *Chandra* exposures toward the Galactic Center (Muno et al., 2003).

5. Conclusion

1. We have modeled populations of short-period semidetached close binary systems containing black holes and low-mass stars under various assumptions about the AML mechanism.

We have found that the "standard" model (based on Skumanich (1972) stellar rotation braking law prescription for AML via magnetic braking) produces unobserved persistent LMBHB, while a model with pure GWR AML produces only transient systems. In the latter model it is possible to reproduce, within uncertainty of observations, the number of LMBHB in the Galaxy, the effective temperatures and masses of the donors in these systems (as inferred from the spectra of the latter). Mass-transfer rates in this model are consistent with the upper limits on $\dot{M}$, if quiescent accretion discs are truncated and, hence, leaky. All this suggests that the strength of MSW AML should be strongly reduced compared to the "standard" model.

In the model without strong MSW it is unnecessary to invoke systems with the donors that are almost at TAMS or already left the latter ($X_c \lesssim 0.01$) in order to explain observed orbital periods of SXT and their mass-exchange rates. More, population of short-period ($P_{\text{orb}} \lesssim 8 - 9$ hr) is formed by systems that overflow Roche lobes unevolved or slightly evolved ($X_c > 0.35$), while evolution of systems with more evolved donors with masses $\simeq 1 M_{\odot}$ opens possibility to explain the origin of LMBHB with longer P_{orb} .

2. An alternative model suggests that if accretion discs are truncated in quiescence they could be cold and stable. Then truncation of discs at radii close to R_{circ} can make discs in virtually all LMBHB cold and stable. In this case SXT outbursts perhaps are not related to DIM and are random events that result from "external factors", like episodes of enhanced mass transfer or variability in truncation radius.

3. Population synthesis models presented above are obtained under assumption of a high value of the product of binding energy parameter of stellar envelopes and expulsion efficiency of common envelopes – $\alpha_{\text{CE}} \times \lambda = 2$. Model with $\alpha_{\text{CE}} \times \lambda = 0.5$ gives similar results (but a reduced total number of systems). Further decrease of $\alpha_{\text{CE}} \times \lambda$ results in

models that are not consistent with observed SXT. The issue of $\alpha_{\text{CE}} \times \lambda$ is still controversial (see, e.g. Tauris and Dewi, 2001; Podsiadlowski et al., 2003) in the sense that there are no strict criteria for defining binding energy of stellar envelopes and there is no clear understanding whether sources other than gravitational energy may contribute to unbinding common envelopes. Apart of common envelopes, crucial role in determining the possibility of formation of LMBHB is played by stellar wind mass-loss in all stages of stellar evolution, since it defines parameters of the system at the beginning of common envelope stage and immediately before supernova explosion that gives birth to black hole. If general notions on mass-loss will not be revised, our results suggest that existence of short-period low-mass X-ray binaries demands high values of $\alpha_{\text{CE}} \times \lambda$ parameter.

6. Acknowledgements

We acknowledge P.P. Eggleton for providing a copy of his evolutionary code, G. Nelemans and G. Dubus for fruitful cooperation and L. Titarchuk and N. Shaposhnikov for useful remarks. LRY acknowledges warm hospitality and support from Institut d'Astrophysique de Paris, Université Pierre et Marie Curie where most of this study was carried out and support by Polish Ministry of Education grant N203 009 31/1466 and Swedish VR grant Dnr 621-2006-3288 which enabled his participation in the conference. LRY is supported by RFBR grant 07-02-00454 and Russian Academy of Sciences Basic Research Program "Origin and Evolution of Stars and Galaxies".

References

- Andronov, N., Pinsonneault, M., Sills, A., Jan. 2003. Cataclysmic Variables: An Empirical Angular Momentum Loss Prescription from Open Cluster Data. *ApJ* 582, 358–368.
- Boissier, S., Prantzos, N., Aug. 1999. Chemo-spectrophotometric evolution of spiral galaxies - I. The model and the Milky Way. *MNRAS* 307, 857–876.
- Charles, P. A., Coe, M. J., Apr. 2006. Optical, ultraviolet and infrared observations of X-ray binaries. In: Lewin, W., van der Klis, M. (Eds.), *Compact stellar X-ray sources*. Cambridge, pp. 215–265.

- Chaty, S., Haswell, C. A., Malzac, J., Hynes, R. I., Shrader, C. R., Cui, W., Dec. 2003. Multiwavelength observations revealing the evolution of the outburst of the black hole XTE J1118+480. *MNRAS* 346, 689–703.
- Chen, W., Shrader, C. R., Livio, M., Dec. 1997. The Properties of X-Ray and Optical Light Curves of X-Ray Novae. *ApJ* 491, 312.
- Collier Cameron, A., Jan. 2002. Magnetic activity in low-mass stars: Do the brakes come off? In: Gänsicke, B. T., Beuermann, K., Reinsch, K. (Eds.), *The Physics of Cataclysmic Variables and Related Objects*. Vol. 261 of *Astronomical Society of the Pacific Conference Series*. p. 11.
- Cox, A. N., 2000. *Allen's astrophysical quantities*, 4th ed. New York: AIP Press; Springer, 2000. Ed. Arthur N. Cox.
- de Kool, M., van den Heuvel, E. P. J., Pylyser, E., 1987. An evolutionary scenario for the black hole binary a0620-00. *A&A* 183, 47–52.
- Done, C., Gierliński, M., Jan. 2006. Truncated disc versus extremely broad iron line in XTE J1650-500. *MNRAS*, 121.
- Dubus, G., Hameury, J.-M., Lasota, J.-P., Jul. 2001. The disc instability model for X-ray transients: Evidence for truncation and irradiation. *A&A* 373, 251–271.
- Dubus, G., Lasota, J., Hameury, J., Charles, P., Feb. 1999. X-ray irradiation in low-mass binary systems. *MNRAS* 303, 139–147.
- Eggleton, P. P., 1971. The evolution of low mass stars. *MNRAS* 151, 351.
- Ergma, E., Fedorova, A., 1998. Evolution of black hole low-mass binaries. *A&A* 338, 69–74.
- Esin, A. A., Lasota, J.-P., Hynes, R. I., Feb. 2000. The 1996 outburst of GRO J1655-40: disc irradiation and enhanced mass transfer. *A&A* 354, 987–994.
- Fryer, C. L., Kalogera, V., Jun. 2001. Theoretical Black Hole Mass Distributions. *ApJ* 554, 548–560.
- Hameury, J. M., Lasota, J. P., King, A. R., Ritter, H., Mar. 1988. Magnetic braking and the evolution of cataclysmic binaries. *MNRAS* 231, 535–547.
- Hobbs, G., Lorimer, D. R., Lyne, A. G., Kramer, M., Jul. 2005. A statistical study of 233 pulsar proper motions. *MNRAS* 360, 974–992.
- Hynes, R. I., Haswell, C. A., Chaty, S., Shrader, C. R., Cui, W., Mar. 2002. The evolving accretion disc in the black hole X-ray transient XTE J1859+226. *MNRAS* 331, 169–179.
- Ivanova, N., Kalogera, V., Jan. 2006. The Brightest Point X-Ray Sources in Elliptical Galaxies and the Mass Spectrum of Accreting Black Holes. *ApJ* 636, 985–994, arXiv:astro-ph/0506471.
- Ivanova, N., Taam, R. E., Dec. 2003. Magnetic Braking Revisited. *ApJ* 599, 516–521.
- King, A. R., Kolb, U., Burderi, L., Jun. 1996. Black hole binaries and x-ray transients. *ApJ* 464, L127.
- Kolb, U., Jun. 1998. Soft x-ray transients in the hertzsprung gap. *MNRAS* 297, 419–426.
- Kroupa, P., Tout, C. A., Gilmore, G., Jun. 1993. The distribution of low-mass stars in the Galactic disc. *MNRAS* 262, 545–587.
- Landau, L. D., Lifshitz, E. M., 1971. "Classical theory of fields", 3rd Edition. Pergamon, Oxford.
- Lasota, J. P., 1996. Mechanisms for Dwarf Nova Outbursts and Soft X-ray Transients (A Critical Review). In: *IAU Symp. 165: Compact Stars in Binaries*. p. 43.
- Lasota, J.-P., Jun. 2001. The disc instability model of dwarf novae and low-mass X-ray binary transients. *New Astronomy Review* 45, 449–508.
- Lasota, J.-P., Dubus, G., Kruk, K., Feb. 2008. Stability of helium accretion discs in ultracompact binaries. ArXiv e-prints 0802.3848.
- Lasota, J.-P., Narayan, R., Yi, I., Oct. 1996. Mechanisms for the outbursts of soft X-ray transients. *A&A* 314, 813–820.
- McClintock, J. E., Remillard, R. A., Apr. 2006. Black hole binaries. In: Lewin, W., van der Klis, M. (Eds.), *Compact stellar X-ray sources*. Cambridge, pp. 157–213.
- Menou, K., Narayan, R., Lasota, J., Mar. 1999. A population of faint nontransient low-mass black hole binaries. *ApJ* 513, 811–826.
- Muno, M. P., Baganoff, F. K., Bautz, M. W., Brandt, W. N., Broos, P. S., Feigelson, E. D., Garmire, G. P., Morris, M. R., Ricker, G. R., Townsley, L. K., May 2003. A Deep Chandra Catalog of X-Ray Point Sources toward the Galactic Center. *ApJ* 589, 225–241.
- Nelemans, G., Yungelson, L. R., Portegies Zwart, S. F., Mar. 2004. Short-period AM CVn systems as optical, X-ray and gravitational-wave sources. *MNRAS* 349, 181–192.
- Nelemans, G., Yungelson, L. R., Portegies Zwart, S. F., Verbunt, F., 2001. Population synthesis for double white dwarfs. i detached systems. *A&A* 365, 491 – 507.
- Orosz, J. A., 2003. Inventory of black hole binaries. In: van der Hucht, K., Herrero, A., Esteban, C. (Eds.), *A Massive Star Odyssey: From Main Sequence to Supernova*. Vol. 212 of *IAU Symposium*.

- p. 365.
- Orosz, J. A., McClintock, J. E., Remillard, R. A., Corbel, S., Nov. 2004. Orbital Parameters for the Black Hole Binary XTE J1650-500. *ApJ* 616, 376–382.
- Paczyński, B., Jan. 1990. A test of the galactic origin of gamma-ray bursts. *ApJ* 348, 485–494.
- Podsiadlowski, P., Rappaport, S., Han, Z., May 2003. On the formation and evolution of black hole binaries. *MNRAS* 341, 385–404.
- Portegies Zwart, S. F., Verbunt, F., 1996. Population synthesis of high-mass binaries. *A&A* 309, 179–196.
- Portegies Zwart, S. F., Yungelson, L. R., 1998. Formation and evolution of binary neutron stars. *A&A* 332, 173.
- Remillard, R. A., McClintock, J. E., Sep. 2006. X-Ray Properties of Black-Hole Binaries. *Ann. Rev. Astron. and Astrophys.* 44, 49–92, arXiv:astro-ph/0606352.
- Romani, R. W., May 1998. A census of low mass black hole binaries. *A&A* 333, 583–590.
- Shakura, N. I., Sunyaev, R. A., 1973. Black holes in binary systems. Observational appearance. *A&A* 24, 337–355.
- Skumanich, A., Feb. 1972. Time Scales for CA II Emission Decay, Rotational Braking, and Lithium Depletion. *ApJ* 171, 565.
- Tauris, T., Dewi, J. D. M., Apr. 2001. On the binding energy parameter of common envelope evolution. *A&A* 369, 170–173.
- van Paradijs, J., Jun. 1996. On the Accretion Instability in Soft X-Ray Transients. *ApJ* 464, L139.
- Verbunt, F., Zwaan, C., Jul. 1981. Magnetic braking in low-mass x-ray binaries. *A&A* 100, L7–L9.
- Webbink, R. F., 1984. Double white dwarfs as progenitors of r coronae borealis stars and type i supernovae. *ApJ* 277, 355.
- White, N. E., van Paradijs, J., 1996. The galactic distribution of black hole candidates in low-mass x-ray binary systems. *ApJ* 473, L25–L29.
- Yungelson, L. R., Lasota, J.-P., Nelemans, G., Dubus, G., van den Heuvel, E. P. J., Dewi, J., Portegies Zwart, S., Aug. 2006. The origin and fate of short-period low-mass black-hole binaries. *A&A* 454, 559–569.